Hydrobiologia

The International Journal on Limnology and Marine Sciences

Contents

Volume 379

Dynamics of abiotic parameters, solute removal and sediment retention in summer-dry headwater streams of western Oregon
by M. Dieterich & N.H. Anderson 1–15

Troglochaetus beranecki Delachaux (Polychaeta, Archiannelida) in esker groundwaters of Finland: a new class of limnic animals for northern Europe
by J. Särkkä & J. Mäkelä 17–21

A new species of Ancorabolidae (Copepoda, Harpacticoida) from the Beagle channel (Chile)
by K.H. George 23–32

Ecology of aquatic Lepidoptera (Crambidae: Nymphulinae) in South Carolina, USA
by C.A. Stoops, P.H. Adler & J.W. McCreadie 33–40

Seasonal coupling of a tropical mangrove forest and an estuarine water column: enhancement of aquatic primary productivity
by V.H. Rivera-Monroy, C.J. Madden, J.W. Day, Jr., R.R. Twilley, F.Vera-Herrera & H. Alvarez-Guillén 41–53

The effect of solar UVA and UVB on subarctic *Daphnia pulicaria* in its natural habitat
by I.D. Zellmer 55–62

Factors influencing the distribution and abundance of diaptomid copepods in high-elevation lakes in the Pacific Northwest, USA
by W.J. Liss, G.L. Larson, E.A. Deimling, L.M. Ganio, R.L. Hoffman & G.A. Lomnicky 63–75

Species of the planktonic diatom genus *Pseudo-nitzschia* of the Pacific coasts of Mexico
by D.U. Hernández-Becerril 77–84

A new species of *Hemicyclops* (Copepoda, Poecilostomatoida) from burrows of the ocypodid crab *Macrophthalmus japonicus* in an estuarine mud-flat in Tokyo Bay, Japan
by H. Itoh & S. Nishida 85–92

The freshwater crab fauna (Crustacea: Brachyura) of the Philippines. V. On a new genus and species of potamid from Palawan island, Philippines
by S.H. Tan & P.K.L. Ng 93–96

Nematodes from the Strait of Magellan and the Beagle Channel (Chile): description of four new species of the Comesomatidae
by G. Chen & M. Vincx 97–110

Stormwater impact to the freshwater Savannas Preserve marsh, Florida, USA
by G.A. Graves, D.G. Strom & B.E. Robson 111–122

Free-living marine nematodes of shrimp culture ponds in Thailand. I. New species of the genera *Diplolaimella* and *Thalassomonhystera* (Monhysteridae) and *Theristus* (Xyalidae)
by K. Kito & C. Aryuthaka 123–133

Effects of acidic pH on benthic macroinvertebrate communities in stream microcosms
by L.A. Courtney & W.H. Clements 135–145
The use of structural dynamic models to explain successes and failures of biomanipulation
by S.E. Jørgensen & R. de Bernardi 147–158
Redescription and taxonomic status of *Sinodiaptomus valkanovi*, a common limnoplanktonic calanoid copepod in Japan, with comparison to the closely related *S. sarsi*
by H. Ueda & S. Ohtsuka 159–168
Interaction between humic substances and carbohydrates in a Pampasic pond (Argentina)
by V.H. Conzonno & A.F. Cirelli 169–174
Relationships between environmental variables and the abundance of cladocerans and copepods in the Atchafalaya River Basin
by N.L. Davidson, Jr., W.E. Kelso & D.A. Rutherford 175–181
Bythotrephes longimanus in the Commonwealth of Independent States: variability, distribution and ecology
by I.A. Grigorovich, O.V. Pashkova, Y.F. Gromova & C.D.A. van Overdijk 183–198
The influence of food quality on the phototactic behaviour of *Daphnia magna* Straus
by E. Michels & L. De Meester 199–206

Notes
Injury and short term mortality of benthic stream fishes – a comparison of collection techniques
by S.J. Cooke, C.M. Bunt & R.S. McKinley 207–211
Troglochaetus beranecki Delachaux (Annelida, Polychaeta): collecting methods and microscopy techniques for SEM and *in vivo* observations
by M. Ivano, S. Manuela & M. Marisa 213–216

Author Index 217